

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representation of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY

As rescanning documents *will not* correct images, please do not report the images to the Image Problem Mailbox.

(51) International Patent Classification ⁵ : C12N 15/31, 15/82, 15/67 A01H 5/00	A1	(11) International Publication Number: WO 91/16432 (43) International Publication Date: 31 October 1991 (31.10.91)
(21) International Application Number: PCT/EP91/00733 (22) International Filing Date: 17 April 1991 (17.04.91) (30) Priority data: 90401055.0 18 April 1990 (18.04.90) EP (34) Countries for which the regional or international application was filed: GB et al. (71) Applicant (for all designated States except US): PLANT GENETIC SYSTEMS N.V. [BE/BE]; Kolonel Bourgstraat 106, B-1040 Brussels (BE). (72) Inventors; and (75) Inventors/Applicants (for US only) : CORNELISSEN, Marc [NL/BE]; Ellebogten 38, B-9070 Heusden (BE). SOËTART, Piet [BE/BE]; Eekhoekstraat 42 A, B-9270 Laarne (BE). STAM, Maike [NL/NL]; Uilenstede 88-576, NL-1183 AL Amstelveen (NL). DOCKX, Jan [BE/BE]; Tentoonstellingslaan 107, B-9000 Gent (BE).		(74) Agents: GUTMANN, Ernest et al.; Ernest Gutmann-Yves Plasseraud S.A., 67, boulevard Haussmann, F-75008 Paris (FR). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i>
(54) Title: MODIFIED BACILLUS THURINGIENSIS INSECTICIDAL-CRYSTAL PROTEIN GENES AND THEIR EXPRESSION IN PLANT CELLS (57) Abstract <p>A DNA fragment, encoding all or an insecticidally-effective part of a Bt crystal protein, is modified by changing A and T sequences to corresponding G and C sequences encoding the same amino acids.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

MODIFIED BACILLUS THURINGIENSIS
INSECTICIDAL-CRYSTAL PROTEIN GENES AND THEIR
EXPRESSION IN PLANT CELLS

5 This invention provides a modified Bacillus
thuringiensis ("Bt") gene (the "modified BtICP gene")
encoding all or an insecticidally-effective portion of
a Bt insecticidal crystal protein ("ICP"). A plant,
transformed with the modified Bt ICP gene can show
10 higher expression levels of the encoded ICP and
improved insect-resistance.

Background of the Invention

15 Plant genetic engineering technology has made
significant progress during the last 10 years. It has
become possible to introduce stably foreign genes into
plants. This has provided exciting opportunities for
modern agriculture. Derivatives of the Ti-plasmid of
the plant pathogen, Agrobacterium tumefaciens, have
20 proven to be efficient and highly versatile vehicles
for the introduction of foreign genes into plants and
plant cells. In addition, a variety of free DNA
delivery methods, such as electroporation,
microinjection, pollen-mediated gene transfer and
25 particle gun technology, have been developed for the
same purpose.

30 The major aim of plant transformations by genetic
engineering has been crop improvement. In an initial
phase, research has been focused on the engineering
into plants of useful traits such as insect-resistance.
In this respect, progress in engineering insect
resistance in transgenic plants has been obtained
through the use of genes, encoding ICPs, from Bt
strains (Vaech t al., 1987). A Bt strain is a sp re
forming gram-positive bacterium that pr duces a

parasp ral crystal which is composed f crystal
prot ins which are specifically toxic against ins ct
larvae. Bt ICPs possess a specific insecticidal
spectrum and display no toxicity towards other animals
5 and humans (Gasser and Fraley, 1989). Therefore, the Bt
ICP genes are highly suited for plant engineering
purposes.

For more than 20 years, Bt crystal spore
preparations have been used as biological insecticides.
10 The commercial use of Bt sprays has however been
limited by high production costs and the instability of
crystal proteins when exposed in the field (Vaech et
al., 1987). The heterogeneity of Bt strains has been
well documented. Strains active against Lepidoptera
15 (Dulmage et al., 1981), Diptera (Goldberg and Margalit,
1977) and Coleoptera (Krieg et al., 1983) have been
described.

Bt strains produce endogenous crystals upon
sporulation. Upon ingestion by insect larvae, the
20 crystals are solubilized in the alkaline environment of
the insect midgut giving rise to a protoxin which is
subsequently proteolytically converted into a toxic
core fragment or toxin of 60-70 kDa. The toxin causes
cytolysis of the epithelial midgut cells. Th
25 specificity of Bt ICPs can be determined by their
interaction with high-affinity binding sites present on
insects' midgut epithelia.

The identification of Bt ICPs and the cloning and
sequencing of Bt ICP genes has been reviewed by Höfte
30 and Whiteley (1989). The Bt ICP genes share a number of
common properties. They generally encode insecticidal
proteins of 130 kDa to 140 kDa or of about 70 kDa,
which contain t xic fragm nts of 60 ± 10 kDa (Höft and
Whiteley, 1989). The Bt ICP gen s have b n classifi d
into four maj r groups according to b th their

structural similarities and insecticidal spectra (Höft and Whiteley, 1989): Lepidoptera-specific (CryI), Lepidoptera- and Diptera-specific (CryII), Coleoptera-specific (CryIII) and Diptera-specific (Cry IV) genes.

5 The Lepidoptera-specific genes (CryI) all encode 130-140 kDa proteins. These proteins are generally synthesized as protoxins. The toxic domain is localized in the N-terminal half of the protoxin. Deletion analysis of several CryI genes confirm that 3' portions

10 of the protoxins are not absolutely required for toxic activity (Schnepp et al., 1989). Cry II genes encode 65 kDa proteins (Widner and Whiteley, 1985). The Cry II A proteins are toxic against both Lepidoptera and Diptera while the Cry II B proteins are toxic only to

15 Lepidopteran insects. The Coleoptera-specific genes (Cry III) generally encode proteins with a molecular weight of about 70 kDa. (Whiteley and Höfte, 1989). The corresponding gene (cry III A) expressed in E. coli directs the synthesis of a 72 kDa protein which is

20 toxic for the Colorado potato beetle. This 72 kDa protein is processed to a 66 kDa protein by spore-associated bacterial proteases which remove the first 57 N-terminal amino acids (Mc Pherson et al., 1988). Deletion analysis demonstrated that this type of gene

25 cannot be truncated at its 3'-end without the loss of toxic activity (Höfte and Whiteley, 1989). Recently, an anti-coleopteran strain, which produces a 130 kDa, protein has also been described (European patent application ("EPA") 89400428.2). The cry IV class of

30 crystal protein genes is composed of a heterogenous group of Diptera-specific crystal protein genes (Höfte and Whiteley, 1989).

The feasibility of generating insect-resistant transgenic crops by using Bt ICPs has been demonstrated. (Vack et al., 1987 ; Fischhoff et al.,

1987 and Barton et al., 1987). Transgenic plants offer an attractive alternative and provide an entirely new approach to insect control in agriculture which is at the same time safe, environmentally attractive and cost-effective. (Meeusen and Warren, 1989). Successful insect control has been observed under field conditions (Delannay et al., 1989 ; Meeusen and Warren, 1989).

In all cases, Agrobacterium-mediated gene transfer has been used to express chimaeric Bt ICP genes in plants (Vaeck et al., 1987; Barton et al., 1987; ~~Fischhoff et al., 1987~~). ~~Bt ICP genes were placed under~~ the control of a strong promoter capable of directing gene expression in plant cells. It is however remarkable that expression levels in plant cells were high enough only to obtain insect-killing levels of Bt ICP genes when truncated genes were used (Vaeck et al., 1987; Barton et al., 1987). None of the transgenic plants containing a full-length Bt ICP gene produced insect-killing activity. Moreover, Barton et al. (1987) showed that tobacco calli transformed with the entire Bt ICP coding sequence became necrotic and died. These results indicate that the Bt ICP gene presents unusual problems that must be overcome to obtain significant levels of expression in plants. Even, when using a truncated Bt ICP gene for plant transformation, the steady state levels of Bt ICP mRNA obtained in transgenic plants are very low relative to levels produced by both an adjacent NPT II-gene, used as a marker, and by other chimeric genes (Barton et al., 1987; Vaeck et al., 1987). Moreover, the Bt ICP mRNA cannot be detected by northern blot analysis. Similar observations were made by Fischhoff et al. (1987); they reported that the level of Bt ICP mRNA was much lower than expected for a chimeric gene expressed from the CaMV35S promoter. In other words, the cytoplasmic

accumulation of the bt mRNA, and consequently the synthesis, the accumulation and thereby the expression of the Bt ICP protein in plant cells, are extremely inefficient. By contrast, in microorganisms, it has
5 been shown that truncated Bt ICP genes are less favorable than full-length genes (Adang et al., 1985), indicating that the inefficient expression is solely related to the heterologous expression of Bt ICP genes in plants.

10 The problem of obtaining significant Bt ICP expression levels in plant cells seems to be inherent and intrinsic to the Bt ICP genes. Furthermore, the relatively low and poor expression levels obtained in plants appears to be a common phenomenon for all Bt ICP
15 genes.

It is known that there are six steps at which gene expression can be controlled in eucaryotes (Darnell, 1982):

- 20 1) Transcriptional control
2) RNA processing control
3) RNA transport control
4) mRNA degradation control
5) translational control
25 6) protein activity control

For all genes, transcriptional control is considered to be of paramount importance (The Molecular Biology of the Cell, 1989).

30 In European patent publications ("EP") 385,962 and 359,472, efforts to modify the codon usage of Bt ICP genes to improve their expression in plant cells have been reported. However, wholesale (i.e., non-selective) changes in codon usage can introduce cryptic regulatory signals in a gene, thereby causing problems in one or

more of the six steps mentioned above for gene expression, and thus inhibiting or interfering with transcription and/or translation of the modified foreign gene in plant cells. For example, changes in
5 codon usage can cause differential rates of mRNA production, producing instability in the mRNA, so produced (e.g., by exposure of regions of the mRNA, unprotected by ribosomes, to attack and degradation by cytoplasmic enzymes). Changes in codon usage also can
10 inadvertently cause inhibition or termination of RNA polymerase II elongation on the so-modified gene.

Summary of the Invention

In accordance with this invention is provided a
15 process for modifying a foreign gene, particularly a Bt ICP gene, whose level and/or rate of expression in plant cells, transformed with the gene, is limited by the rate and/or level of nuclear production of an mRNA encoded by the gene; the process comprises the step of
20 changing adenine and thymine sequences to corresponding guanine and cytosine sequences encoding the same amino acids in a plurality of translational codons of the gene that would otherwise directly or indirectly cause a nuclear event which would negatively control (i.e.,
25 inhibit or interfere with) transcription, nuclear accumulation and/or nuclear export of the mRNA, particularly transcription, quite particularly elongation of transcription by RNA polymerase II of the plant cells. Preferably, the adenine and thymine
30 sequences are changed to cytosine and guanine sequences in translational codons of at least one region of the gene which, during transcription, would otherwise have thereon a relatively low percentage of RNA polymerase II as compared to an other adjacent upstream (i.e., 5') region of the gene.

Also in accordance with this invention is provided the modified Bt ICP gene resulting from the process.

Further in accordance with this invention, a process is provided for improving the resistance of a plant against insect pests by transforming the plant cell genome with at least one modified Bt ICP gene.

This invention also relates to a chimaeric gene that can be used to transform plant cells and that contains the following operably-linked DNA fragments in the same transcriptional unit:

- 1) the modified Bt ICP gene;
- 2) a promoter suitable for directing transcription of the modified Bt ICP gene in the plant cells;
- and
- 3) suitable transcript 3' end formation and polyadenylation signals for expressing the modified Bt ICP gene in the plant cells.

This invention further relates to:

- a cell of a plant, the nuclear genome of which has been transformed to contain, preferably stably integrated therein, the modified Bt ICP gene, particularly the chimaeric gene;
- cell cultures consisting of the plant cell;
- a plant which is regenerated from the transformed plant cell or is produced from the so-regenerated plant, the genome of which contains the modified Bt ICP gene, particularly the chimaeric gene, and which shows improved resistance to insect pests;
- seeds of the plant; and
- a vector for stably transforming the nuclear genome of plant cells with the modified Bt ICP gene, particularly the chimaeric gene.

Detailed Description of the Invention

As used herein, "Bt ICP" should be understood as an intact protein or a part thereof which has insecticidal activity and which can be produced in nature by B. thuringiensis. A Bt ICP can be a protoxin, as well as an active toxin or other insecticidal truncated part of a protoxin which need not be crystalline and which need not be a naturally occurring protein. An example of a Bt ICP is a Bt2 insecticidal crystal protein (Höfte et al., 1986), as well as its insecticidally effective parts which are truncated at its C- and/or N-terminal ends towards its trypsin cleavage site(s) and preferably having a molecular weight of 60-80 kDa. Other examples of Bt ICPs are: Bt2, Bt3, Bt4, Bt13, Bt14, Bt15, Bt18, Bt21, Bt22, Bt73, Bt208, Bt245, BtI260 and BtI109P as disclosed in PCT publications WO90/15139 and WO90/09445, in Höfte and Whiteley (1989) and in EPA 90403724.9.

As used herein, "protoxin" should be understood as the primary translation product of a full-length gene encoding a Bt ICP.

As used herein, "toxin" or "active toxin" or "toxic core" should all be understood as a part of a protoxin which can be obtained by protease (e.g., by trypsin) cleavage and has insecticidal activity.

As used herein, "truncated Bt gene" should be understood as a fragment of a full-length Bt gene which still encodes at least the toxic part of the Bt ICP, preferentially the toxin.

As used herein, "modified Bt ICP gene" should be understood as a DNA sequence which encodes a Bt ICP, and in which the content of adenine ("A") and thymine ("T") has been changed to guanine ("G") and cytosine ("C") in codons, preferably at least 3, in at least one region of the DNA sequence without affecting the

original amino acid sequence of the Bt ICP. Preferably in at least two regions, especially in at least three regions, of the DNA sequence, the A and T content is changed to G and C in at least 3 codons. For regions
5 downstream of the translation initiation site of the DNA sequence, it is preferred that the A-T content of at least about 10 codons, particularly at least about 33 codons, be changed to G-C.

By "region" of a modified Bt ICP gene is meant any
10 sequence encoding at least three translational codons which affect expression of the gene in plants.

In accordance with this invention, it has been shown by means of mRNA turn-over studies that the expression pathway of a Bt ICP gene, such as bt2, bt14,
15 bt15 and bt18, is specifically inhibited at the nuclear level in plant cells. In a further analysis, nuclei of transgenic tobacco plants, i.e., N28 - 220 (Vaeck et al., 1987), were used in a nuclear run-on assay to determine the distribution and the relative efficiency
20 of RNA polymerase II complexes to initiate transcription of chimaeric Bt ICP plant genes. In this regard, the run-on assay has been used to determine initially the relative efficiency of RNA polymerase II complexes to initiate transcription of Bt ICP genes and
25 thereafter to determine the relative distribution and migration efficiency of the RNA polymerase II complexes on the Bt ICP genes.

N28 - 220 contains the bt884 fragment under control of the TR 2' promoter as a chimaeric gene.
30 Bt884 is a 5' fragment of the bt2 gene (Höfte et al., 1986) up to codon 610 (Vaeck et al., 1987). Using nuclear run-on analysis, isolated nuclei of N28 - 220 were incubated with highly labeled radioactive RNA precursors, so that the RNA transcripts being synthesized at the time became radioactively labeled.

Th RNA polymeras II molecules caught in th act of transcripti n in the cell continue el ngating the same RNA molecules in vitro.

5 The nuclear run-on assays of nuclei of N28 - 220 culture (non-induced cells and induced cells, TR1'-neo, TR2'-bt884) revealed that transcription from the TR1' and TR2' promoters is about equally efficient. This implies that the low Bt ICP (i.e., Bt884) expression levels are not due to a specifically reduced
10 transcriptional activity of the TR2' promoter. However, nuclear run-on analysis with N28 - 220 nuclei indicated that transcription elongation of the nascent Bt ICP mRNA is impaired somewhere between 700 to 1000 nucleotides downstream of the start of transcription.
15 This means that RNA polymerase II is not able to transcribe the Bt ICP coding sequence with 100 % efficiency. Filter binding assays using labeled Bt DNA fragments spanning this region and protein extract prepared from tobacco nuclei reveal that this DNA
20 region undergoes specific interactions with proteins present in nuclei. These interactions are the prime candidates that cause or affect the impaired elongation of transcription by RNA polymerase II through this region. By modification of this region to abolish
25 specific protein binding, Bt ICP expression levels will increase. However, other mechanisms responsible for impaired elongation in this region cannot be excluded.

Further in accordance with this invention, sequences within the coding region involved in negative
30 control of cytoplasmic Bt ICP mRNA levels have been identified by deletion analysis. To this end, 24 deletion derivatives of pVE36 have been constructed. Three main typ s of del tion mutants have been c nstructed (se fig. 3):

- 5' end deletions
- 3' end deletions
- internal deletions.

5 The expression of a mutant hybrid bt2-neo gene (encoding a fusion protein of Bt2 (Höfte et al., 1986) and NPTII) has been studied by means of transient expression experiments using the cat gene as a reference. To this end, the neo mRNA levels were
10 measured in relation to cat mRNA levels in RNA extracts of SR1 protoplasts. The ratio between the neo and cat mRNA level was used to quantify on a relative basis the nptII transcript (i.e., mRNA) levels produced by the different constructions. These experiments show that
15 progressive deletions of the carboxy-terminal (i. e., 3') part or the amino-terminal (i. e., 5') part of the Bt ICP coding sequence result in a gradual increase of the nptII transcript level. Furthermore, since the changes in transcript levels are not very abrupt, these
20 results suggest that the low transcript levels produced by Bt ICP genes are not controlled by a single factor. Nevertheless, individual modifications of bt2 coding sequence can significantly reduce the interference and/or inhibition of the expression of the mRNA encoded
25 by Bt ICP genes in plant cells at the level of transcript elongation, nuclear accumulation and nuclear export. The modification(s) may also affect cytoplasmic regulation and metabolism of such mRNAs and their translation.

30 Deletion analysis clearly indicates that several internal sequences, located within the Bt ICP coding region, might be involved in the negative regulation of the Bt ICP expression. By way of example, a 326 bp region (fig. 6b) was identified in the bt2 gene that is involved in the negative control of BT ICP expression

and that is located between nucleotide position 674 and nucleotide position 1000, particularly a 268 bp region between nucleotide positions 733 and 1000, quite particularly a 29 bp region between nucleotide positions 765 and 794 which carries two perfect CCAAT boxes which are known to be able to cause a reduction in elongation efficiency and termination of transcription by RNA polymerase II in animal systems (Connelly and Manley, 1989). This internal gene fragment or inhibitory zone may itself comprise a plurality of inhibitory zones which reduce Bt ICP expression levels or which interact directly or indirectly with other zones to inhibit or interfere with expression. Codon usage of this inhibitory zone has been modified in a second step by substituting A - T with G - C without affecting the amino acid sequence. In this regard, this internal 326 bp fragment (fig. 6b) has been replaced with a modified Bt ICP fragment of this invention containing 59 modified codons. The effect of such modification of this inhibitory zone on Bt ICP expression has been analyzed both in transient and stable plant transformants. The results show that such modification of codon usage causes a significant increase of Bt ICP expression levels and hence improved insect-resistance.

In addition, N-terminal deletion mutants of the bt2 gene have been made by deleting the first N-terminal 28 amino acids (Höfte et al., 1986). It is known for the bt2 gene that the first 28 codons can be deleted without loss of toxicity (Höfte et al., 1986; Vaeck et al., 1987). Also, codon usage for three codons, 29 to 31, has been changed in accordance with this invention by replacing A - T with G - C without affecting the amino acid sequence. Furthermore, an optimal translation initiation (ATG) site was created

based on the consensus sequence of Joshi (1987) as shown in fig. 6a. Plants transformed with this modified Bt ICP gene show significantly higher Bt ICP expression levels.

5 In accordance with this invention, all or part of a modified Bt ICP gene of the invention can be stably inserted in a conventional manner into the nuclear genome of a plant cell, and the so-transformed plant cell can be used to produce a transgenic plant showing
10 improved expression of the Bt ICP gene. In this regard, a disarmed Ti-plasmid, containing the modified Bt ICP gene, in Agrobacterium (e.g., A. tumefaciens) can be used to transform a plant cell using the procedures described, for example, in EP 116,718 and EP 270,822,
15 PCT publication 84/02913, EPA 87400544.0 and Gould et al. (1991) (which are incorporated herein by reference). Preferred Ti-plasmid vectors contain the foreign DNA sequence between the border sequence, or at least located to the left of the right border sequence,
20 of the T-DNA of the Ti-plasmid. Of course, other types of vectors can be used to transform the plant cell, using procedures such as direct gene transfer (as described, for example, in EP 233,247), pollen mediated transformation (as described, for example, in EP
25 270,356, PCT publication WO 85/01856, and US patent 4,684,611), plant RNA virus-mediated transformation (as described, for example, in EP 67,553 and US patent 4,407,956), liposome-mediated transformation (as described, for example, in US patent 4,536,475) and
30 other methods such as the recently described methods for transforming certain lines of corn (Fromm et al., 1990; Gordon-Kamm et al., 1990).

Preferably, the modified Bt ICP gene is inserted in a plant genome downstream of, and under the control of, a promoter which can direct the expression of the

g n in the plant cells. Preferred promoters include, but are not limited to, the strong constitutive 35S promoter (Odell et al., 1985) of cauliflower mosaic virus; 35S promoters have been obtained from different
5 isolates (Hull and Howell, Virology 86, 482-493 (1987)). Other preferred promoters include the TR1' promoter and the TR2' promoter (Velten et al., 1984). Alternatively, a promoter can be utilized which is not constitutive but rather is specific for one or more
10 tissues or organs. For example, the modified Bt ICP gene can be selectively expressed in the green tissues of a plant by placing the gene under the control of a light-inducible promoter such as the promoter of the ribulose - 1,5 - phosphate - carboxylase small subunit
15 gene as described in EPA 86300291.1. Another alternative is to use a promoter whose expression is inducible by temperature or chemical factors.

It is also preferred that the modified Bt ICP gene be inserted upstream of suitable 3' transcription
20 regulation signals (i.e., transcript 3' end formation and polyadenylation signals) such as the 3' untranslated end of the octopine synthase gene (Gielen et al., 1984) or T-DNA gene .7 (Velten and Schell, 1985).

25 The resulting transformed plant of this invention shows improved expression of the modified Bt ICP gene and hence is characterized by the production of high levels of Bt ICP. Such a plant can be used in a conventional breeding scheme to produce more
30 transformed plants with the same improved insect-resistance characteristics or to introduce the modified Bt ICP gene into other varieties of the same or related plant species. Seeds, which are obtained from the transformed plants, contain the modified BtICP gene as a stable genomic insert.

Furthermore, at least two modified BtICP genes, coding for two non-competitively binding anti-Lepidopteran or anti-Coleopteran Bt ICPs, can be cloned into a plant expression vector (EPA 89401499.2). Plants transformed with such a vector are characterized by the simultaneous expression of at least two modified BtICP genes. The resulting transgenic plant is particularly useful to prevent or delay development of resistance to Bt ICP of insects feeding on the plant.

The following Examples illustrate the invention and are not intended to limit its scope. The Figures, referred to in the Examples, are as follows:

Fig.1 -- Comparison of the transcription initiation frequency of RNA polymerase II complexes in nuclei of N28-220. Hybridisation efficiencies of labeled nptII mRNA and Bt ICP mRNA with their complementary DNA counterparts present on a Southern blot were compared. DNA fragments were obtained from a digest of plasmid pGSH163. A schematic view of the region is given. The lengths of the fragments blotted on Hybond-N filter (1), the homologous genes on plasmid pGSH163 (2), and the densitometric values (3) are as follows:

Digest:	1	2	3
BamHI/HindIII	2358	<u>neo</u>	12386
	1695	<u>bt2</u>	6565
	154	<u>bt2</u>	-
	6250	vector	-

Fig. 2a -- Determination of the distribution of the RNA polymerase II complexes on the Bt ICP coding sequence in nuclei of N28-220. The hybridisation of labeled RNA prepared by nuclear run on with DNA fragments of the Bt ICP coding sequence was quantitated. The restriction

fragments and scanning values are given in the table and figure. The scanning value is proportional to "X", the size of the DNA fragment and the # UTP per RNA fragment hybridising. "X" is directly proportional to the number of RNA polymerases passing through the DNA fragment. "X" is proportional to the scanning value divided by the number of UTPs. The X values of the different restriction fragments are shown in the figure. In this regard, conversion of the different densitometric values into relative hybridisation efficiencies by normalising the values of the number of dATPs present in the DNA fragment, complementary to the hybridising RNA, generates the value "X". "X" is a relative measure of the number and the length of the extension of the transcripts. "X" thus reflects the number of RNA polymerases transcribing a specific DNA sequence and their elongation rate. DNA fragments present on the Southern digests of plasmid DNA of plant vector pGSH163 each have the following lengths of fragments blotted on Hybond-N filter (1), homologous genes on plasmid pGSH163 (2) and densitometric values (3):

Digest:	1	2	3
25 BamHI/EcoRI	8877	<u>neo</u>	15333
	726	<u>bt2</u> (2)	2926
	583	<u>bt2</u> (3)	635
	271	<u>bt2</u> (1)	-
30 BamHI/EcoRV	8887	<u>neo</u>	15182
	84	<u>bt2</u>	2466
	729	<u>bt2</u>	1102
BamHI/HindIII	6250	-	-
	2358	<u>neo</u>	12386

17

		1695	<u>bt2</u>	6565
		154	-	-
	BamHI/SacI	8053	<u>neo</u>	14194
5		1353	<u>bt2(1)</u>	4572
		1051	<u>bt2(2)</u>	615
	XmnI	4973	<u>neo</u>	13219
		2107	-	-
		1401	-	-
10		729	<u>bt2(3)</u>	736
		628	<u>bt2(2)</u>	1817
		305	<u>bt2(4)</u>	-
		188	<u>bt2(5)</u>	-
15		120	<u>bt2(1)</u>	-

Fig. 2b -- Schematic view of nine bt884 DNA fragments that were inserted into the polylinker of M13 vectors, MP18 and MP19 (Yanisch-Perron et al., 1985). The Bt ICP coding sequence is shown from AUG to 1600 nucleotides downstream. The relevant restriction sites and sizes of the DNA fragments are indicated. The nucleotide numbering is relative to the AUG. The subclones were named pJD71, pJD72, pJD73, etc. (to pJD79), as indicated. The inserts were oriented into the M13 vector such that single stranded M13 carried the fragments of the Bt ICP coding sequence in an anti-sense orientation.

Fig. 2c -- Schematic representation of three nuclear run-on analyses with N28-220 nuclei as described by Cox and Goldberg (1988). Assays were performed for periods of 5, 10 and 30 minutes. The labeled nuclear RNA was allowed to hybridize with 5 µg of single stranded pJD71-pJD79 and MP18 DNA, which were immobilized on nylon membranes. The membranes were autoradiographed,

and densitometric values were obtained by scanning the autoradiographs. The abscissa shows the nucleotide position relative to the AUG of the Bt (i.e., bt2) coding sequence. The center of each of the single stranded Bt DNA fragments is indicated in the graph. The ordinate gives the relative hybridisation signal for each fragment corrected for the number of dATPs in the fragment and adjusted to 100% for the value of pJD71 for each of the three incubation periods. All values are corrected for non-specific hybridisation to single stranded HP10 DNA. The relative values are a measure for the reactivation of bt mRNA synthesis by RNA polymerase II. The assay does not distinguish between the number of mRNA extensions and the length of mRNA extensions.

Fig. 3 -- Construction of deletion mutants of the bt860-neo gene to measure the effect on cytoplasmic Bt ICP mRNA levels. The parental vector pVE36 is shown. The following deletion mutants were generated:

1. PJD50: pJD50 was derived from pVE36 by digesting with BamHI and SphI. The 5' and 3' protruding ends were filled in with Klenow DNA polymerase I enzyme. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
2. PJD51: pJD51 was derived from pVE36 by digesting with SpeI and SphI. The 5' and 3' protruding ends were filled in with Klenow DNA polymerase I enzyme. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

3. PJD52: pJD52 was derived from pVE36 by dig sting
with EcoRV and SphI. The 5'and 3'
protruding ends were filled in with Klen w
DNA polymerase I enzyme. The treated DNA
was ligated and then used to transform
MC1061 cells. Transformants were selected
for amp^r phenotype.
4. PJD53: pJD53 was derived from pVE36 by digesting
with XcaI and SphI. The 3' protruding ends
were filled in with Klenow DNA polymerase
I enzyme. The treated DNA was ligated and
then used to transform MC1061 cells.
Transformants were selected for amp^r
phenotype.
5. PJD54: pJD54 was derived from pVE36 by digesting
with AflII and SphI. The 5' and 3'
protruding ends were filled in with Klenow
DNA polymerase I enzyme. The treated DNA
was ligated and then used to transform
MC1061 cells. Transformants were selected
for amp^r phenotype.
6. PJD55: pJD55 was derived from pVE36 by digesting
with ClaI and SphI. The 5' and 3'
protruding ends were filled in with Klenow
DNA polymerase I enzyme. The treated DNA
was ligated and then used to transform
MC1061 cells. Transformants were selected
for amp^r phenotype.
7. PJD56: pJD56 was derived from pVE36 by digesting
with XhoI and SphI. The 5' and 3'
protruding ends were filled in with Klenow
DNA polymerase I enzyme. The treated DNA
was ligated and then used t transform
MC1061 cells. Transf rmants wer sel cted
for amp^r ph notype.

- 5 8. PJD57: pJD57 was derived from pVE36 by digesting with AflIII and BamHI. The 5' and 3' protruding ends were filled in with Klenow DNA polymerase I enzyme. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
- 10 9. PJD58: pJD58 was derived from pVE36 by digesting with XcaI and BamHI. The 5' protruding ends were filled in with Klenow DNA ~~polymerase I enzyme. The treated DNA was~~ ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
- 15 10. PJD59: pJD59 was derived from pVE36 by digesting with EcoRV and BamHI. The 5' protruding ends were filled in with Klenow DNA polymerase I enzyme. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
- 20 11. PJD60: pJD60 was derived from pVE36 by digesting with SpeI and BamHI. The 5' protruding ends were filled in with Klenow DNA polymerase I enzyme. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
- 25 12. PJD61: PJD61 was derived from PJD50. PVE36 was digested with XbaI and filled in with Klenow polymerase I. PJD50 was linearized with BamHI and filled in with Klenow polymerase I. The 375bp XbaI fragment of PVE36 was ligated in the filled in BamHI of PJD50. The ligation mixture was used to
- 30

transform MC1061 cells. Transformants were selected for amp^r phenotype.

5
10
13. PJD62: PJD62 was derived from PJD50. PVE36 was digested with XcaI and EcoRV. PJD50 was linearized with BamHI and filled in with Klenow polymerase I. The 367bp XcaI-EcoRV fragment of PVE36 was ligated in the filled in BamHI of pJD50. The ligation mixture was used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

15
20
14. PJD63: PJD63 was derived from PJD50. PVE36 was digested with XcaI and EcoRV. PJD50 was linearized with BamHI and filled in with Klenow polymerase I. The 474bp XcaI-EcoRV fragment of PVE36 was ligated in the filled in BamHI of pJD50. The ligation mixture was used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

25
30
15. PJD64: PJD64 was derived from PJD50. PVE36 was digested with EcoRI and EcoRV and filled in with Klenow polymerase I. PJD50 was linearized with BamHI and filled in with Klenow polymerase I. The 458bp EcoRI-EcoRV fragment of PVE36 was ligated in the filled in BamHI of pJD50. The ligation mixture was used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

16. PJD65: PJD65 was derived from PJD50. PVE36 was digested with EcoRI and XbaI and filled in with Klenow polymerase I. PJD50 was linearized with BamHI and filled in with Klenow polymerase I. The 327bp EcoRI-XbaI

fragment f PVE36 was ligated in the filled in BamHI of pJD50. The ligation mixture was used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

5

17. PJD66: PJD66 was derived from PJD50. PVE36 was digested with SpeI and XcaI and filled in with Klenow polymerase I. PJD50 was linearized with BamHI and filled in with Klenow polymerase I. The 1021bp SpeI-XcaI

10

~~Fragment of PVE36 was ligated in the~~
filled in BamHI of pJD50. The ligation mixture was used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

15

18. PPS56D1: PPS56D1 was derived from PJD56 by digesting with EcoRV. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

20

19. PPS56D2: PPS56D2 was derived from PJD56 by digesting with XcaI and AflII. The 5' protruding ends were filled in with Klenow polymerase I. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

25

20. PPS56D3: PPS56D3 was derived from PJD56 by digesting with SpeI and EcoRV. The 5' protruding ends were filled in with Klenow polymerase I. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

30

21. PPS56D4: PPS56D4 was derived from PJD56 by digesting with XcaI and partially with EcoRV. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
22. PPS56D6: PPS56D6 was derived from PJD56 by digesting with SpeI and partially with EcoRV. The 5' protruding ends were filled in with Klenow polymerase I. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
23. PPS56D7: PPS56D7 was derived from PJD56 by digesting with SpeI and XcaI. The 5' protruding ends were filled in with Klenow polymerase I. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.
24. PPS56D8: PPS56D8 was derived from PPS56D2 by digesting with SpeI and partially with EcoRV. The 5' protruding ends were filled in with Klenow polymerase I. The treated DNA was ligated and then used to transform MC1061 cells. Transformants were selected for amp^r phenotype.

Fig. 4 -- Effect of deletions in the Bt ICP coding sequence on cytoplasmic Bt ICP mRNA levels. The cytoplasmic mRNA levels specified by the invariable cat reference gene and the different Bt ICP deletion mutants described in fig. 3 are listed in the table. The measurements were converted into relative Bt ICP mRNA abundances. Bt ICP and cat mRNA quantizations were

don as described by Corn liss n (1989). Total RNA was
sl t bl tt d and hybridised with radioactively lab l d
RNA complementary to the neo and cat coding sequences.
Values were quantitated with the aid of calibration
5 curves of cold cat and Bt ICP riboprobe transcripts.

Fig. 5 -- Relative transcript levels produced by the
deletion derivatives of pVE36.

10 Fig. 6a -- Schematic presentation of the synthetic DNA
sequences used to introduce a N-terminal deletion and a
change of the codons 29, 30 and 31 of the bt2 coding
sequence. The oligo nucleotides were annealed according
to Engler et al. (1988) and cloned into the BstXI
restriction site of plasmid pVE36, yielding pPS027. The
15 7360 bp fragment of pPS027 was ligated to the the 1177
bp ClaI restriction fragment of pVE36, yielding plasmid
pPS028. pPS028 is identical to pVE36 apart for the N-
terminal modification.

20 Fig. 6b -- Schematic presentation of the synthetic DNA
sequences used to introduce an internal modification
into the bt2 coding sequence. The oligonucleotides were
annealed and ligated as described by Engler et al.
(1988) and the resulting concatemeric DNA fragment was
25 cut with the restriction enzymes XbaI and EcoRI to
release the modified 327 bp XbaI-EcoRI restriction
fragment. This fragment was ligated into the 3530 bp
EcoRI-XbaI fragment of pPS023 which is a pUC19
derivative (Yanisch-Perron et al., 1985) that carries
30 the 1533 bp AflIII (filled in) BamHI fragment of pVE36
in the HindIII (filled in) BamHI site of pUC19,
resulting in plasmid pPS024. Plasmid pPS024 was
lin aris d by dig sti n with restriction enzym XbaI
and th 375 bp XbaI r stricti n fragm nt of pPS023 was
intr duced resulting in pPS025. Th 1177 bp ClaI

fragment of pPS025 was introduced in the 7360 bp *Cla*I restriction fragment of pPS027 yielding pPS029. pPS029 is identical to pVE36 but carries both the amino-terminal modification and the internal modification of the Bt ICP coding sequence.

Fig. 6c -- Nucleotide sequences 800 to 4000 of the plasmids pVE36 and pPS029. "x" refers to not known nucleotides.

Fig. 7 -- Schematic presentation of the effect of the mutations on the AT content of the Bt ICP plant gene. The modified regions are indicated.

Fig. 8a -- Schematic presentation of the plasmid constructions used in the transient expression assay. The relevant genes are indicated.

Fig. 8b -- Accumulation profiles of CAT (Neumann et al., 1987) and the modified BtICP (Engvall and Pesce, 1978) in a typical transient expression assay.

Unless otherwise stated in the Examples, all procedures for making and manipulating recombinant DNA are carried out by the standardized procedures described in Sambrook et al., Molecular Cloning - A laboratory Manual, Cold Spring Harbor Laboratory (1989).

Example 1. Determination of the Efficiency of Transcription Initiation

The relative efficiency of RNA polymerase II complexes to initiate transcription at chimaeric BtICP plant genes was studied, using transgenic plant N28-220 which is described by Vaeck et al. (1987) and contains copies of the T-DNA of plasmid pGSH163. This T-DNA carries the chimaeric plant genes P_{Tr2} bt8843'g7 and

P_{TR1}neo3'ocs. Nuclei of 25 g f induced leaves of N28-220 were prepared according to Cox and Goldberg (1988) and stored the nuclei at a temperature of -70°C. This method causes the nascent precursor mRNA chains and the RNA polymerase II complexes to halt while the complexes remain associated at the DNA. A batch of these nuclei was assayed for the ability to incorporate radioactively labeled UTP as a measure for the transcriptional viability of the nuclei (Cox and Goldberg (1988). This incorporation could be successfully repressed by addition of ~~cr~~amanitin to a final concentration of 2 µg/ml. This shows that the UTP incorporation was due to transcript elongation by RNA polymerase II and that RNA synthesis on the protein coding genes which are occupied by RNA polymerase II can be reactivated under the appropriate experimental conditions.

Batches of the nuclei of N28-220 were used to synthesize radioactively labeled RNA as described by Cox and Goldberg (1988). The radioactive RNA synthesized is a direct representation of the distribution of the RNA polymerases II complexes on the DNA in the nuclei. As the DNA of N28-220 carries two genes which can be assayed, namely the chimaeric ne gene and the chimaeric Bt ICP gene, it is possible to compare the distribution of RNA polymerase II complexes on these two genes. To this end, the radioactive RNA was extracted from the nuclei according to Cox and Goldberg (1988) and used as a probe in a conventional Southern hybridisation. The Southern blot contained DNA fragments carrying the Bt ICP and neo coding sequences in a molar excess relative to the neo and Bt ICP RNA species present in the radioactive probe. A detailed description of the Southern blot is given in fig. 1. The hybridisation experiment resulted in hybridisation

signals to both the neo and Bt ICP coding sequences (fig. 1). Densitometric scanning showed that the intensity of the hybridisation signal to the neo and Bt ICP coding regions was nearly identical. This result implies that the number of transcripts initiating from the TR dual promoter is about similar in both directions. As in plant N28-220 the cytoplasmic neo mRNA level is several magnitudes higher than that of Bt ICP; this shows that the Bt ICP coding sequence indeed negatively controls accumulation of cytoplasmic Bt ICP mRNA, but that this phenomenon is not due to a dominant negative effect on transcription initiation of the chimaeric Bt ICP plant gene.

15 Example 2. Transcription Elongation

The relative distribution of RNA polymerase II complexes on the Bt ICP plant genes present in transgenic plant N28-220 which is described by Vaecck et al. (1987) was investigated. To this end, a second experiment was carried out with batches of the nuclei of N28-220 described in Example 1.

The nuclei were incubated as described by Cox and Goldberg (1988) to synthesize radioactively labeled RNA. The radioactive RNA was extracted as described previously to provide a probe for a Southern hybridisation. The Southern blot prepared for this experiment contained several fragments of the Bt ICP coding sequence in molar excess relative to the complementary RNA present in the probe. The rationale of the experiment was that if the RNA polymerase II complexes were equally distributed over the Bt ICP coding region, the hybridisation with the different Bt ICP DNA fragments present on the Southern blot would be proportional to the size and dATP content of the different fragments. A detailed description of the DNA

fragments present on the Southern is given in fig 2a. The hybridisation of the radioactive RNA extract from the nuclei of N28-220 with the Southern revealed that the complete Bt ICP coding sequence as present in
5 N28-220 is transcribed by RNA polymerase II.

Quantification of the hybridisation signals by densitometric scanning of the autoradiogram showed that more radioactively labeled RNA was hybridising with DNA fragments representing Bt ICP sequences located 5' on
10 the Bt ICP coding sequence than with Bt ICP sequences located 3' on the Bt ICP coding sequence. The actual values are given in fig 2a. This in vitro experiment demonstrates that in vivo the RNA polymerases are not evenly distributed over the Bt ICP coding sequence.

15 The site(s) involved in reducing the RNA polymerase II elongation were then determined more accurately. Nine M13 derivatives were made that carry overlapping fragments of the Bt2 coding sequence spanning the region from the AUG to 1584 nucleotides
20 downstream. The inserts were oriented into the vector such that, in single stranded M13 derivatives, the Bt sequences were complementary to the Bt transcript. A schematic view of the M13 clones is given in fig. 2b.

25 A molar excess of each single stranded anti-Bt DNA was bound to nylon filters to serve as a DNA target for hybridisation with labeled RNA prepared from nuclear run-on assays with N28-220 nuclei as described by Cox and Goldberg (1988). Three nuclear run-ons that differed only in their time period of incubation were
30 carried out simultaneously. The incubation time determines the length of extension of the nascent mRNA chain. Shorter incubation periods give a more accurate view of the position of the RNA polymerase II complexes relative to the substrate DNA and their ability to elongate at the moment of the start of incubation.

Hence, the shorter the in vitro incubation period, the more accurate the view in predicting the in vivo situation.

5 The results are shown in fig. 2c. The data for the 5 minute incubation show that, in vivo, at a very discrete inhibitory zone along the bt2 coding sequence, one or more factors interfere with transcript elongation and that such factor(s) remain present in such inhibitory zone during the course of the in vitro
10 mRNA extension reaction. Increased incubation periods show that, on a subset of DNA templates, RNA synthesis resumed downstream of such inhibitory zone in this assay without significantly removing the inhibition in the inhibitory zone itself. In this regard, the data
15 indicate that:

1. The inhibitory zone causes the RNA polymerases to pause and not to terminate.
2. This pause is only transitory for a small fraction
20 of the Bt DNA templates which were used.
3. The continued RNA polymerase elongation, downstream of the inhibitory zone, is done by a large number of polymerases on the relatively small fraction of the Bt DNA templates.

25 It is believed, therefore, that low cytoplasmic Bt mRNA levels are due at least in part to inefficient production of precursor mRNA caused by inefficient elongation of a nascent transcript and/or stalling of
30 RNA polymerase II complexes from transcribing at an inhibitory zone.

The inhibitory zone was assayed for its ability to interact with proteins present in nuclei of tobacco protoplasts. A crude nuclear extract was prepared from tobacco SR1 leaf protoplasts according to Luthe and

Quatran (1980) and used for filter binding as ay essentially as described by Diffley and Stillman (1986). 100 ng samples of protein extract were mixed with different amounts of radioactively labeled 532 bp
5 XbaI-AccI bt884 DNA fragment, ranging from 0 to 1670 picomolar, in a final volume of 0.150 ml binding buffer (10 mM Tris pH 7.5, 50 mM NaCl, 1 mM DTT, 1 mM EDTA and 5% glycerol). After 45 minutes incubation at room
10 temperature, the samples were filtered through an alkali-washed nitrocellulose membrane and washed twice ~~with 0.150 ml of an ice-cold solution containing 10 mM~~ Tris pH 7.5, 50 mM NaCl and 1 mM EDTA. The retention of DNA-protein complex was quantified by scintillation counting and revealed that the binding had a
15 dissociation constant in the 100 picomolar range. The binding was not affected by preincubation of the nuclear extract with a molecular excess of a specific competitor DNA.

20 Example 3. Construction of Deletion Mutants

The previous two examples demonstrate that the Bt ICP coding sequence in a chimaeric plant gene negatively affects the cytoplasmic Bt ICP mRNA level directed by the chimaeric plant gene. It is shown that
25 this negative control is not at the level of transcription initiation but at least in part due to a reduced ability of RNA polymerase II to generate precursor Bt ICP mRNA. A deletion analysis of the chimaeric Bt ICP plant gene was performed to identify
30 whether impaired transcription elongation is the exclusive mechanism by which the Bt ICP sequence interferes with gene expression. The rationale of the experiment is that the introduction of specific deletions in the Bt ICP coding region could rem v r inactivate the s quence element(s) r sponsible f r th

negative control. As a result such mutant gene would direct an increased level of cytoplasmic mRNA. This method can therefore be used to map and identify the sequence(s) involved in the negative control.

5 To perform this analysis, a deletion series of the bt860-neo gene (Vaech et al., 1987) was made. Fig. 3 gives a schematic representation. The resultant deletion derivatives do not specify a Bt ICP and therefore are assayed at RNA level only. In order to
10 obtain accurate Bt ICP mRNA concentration values, the deletion mutants were compared in a transient expression system using tobacco leaf protoplasts of SR1 (Cornelissen and Vandewiele, 1989). The relative mRNA abundances were calculated using a correction factor
15 provided by the mRNA level specified by the cat reference gene present on the same plasmid as the mutant Bt ICP gene. Four hours after introduction of the genes the tobacco leaf protoplasts were harvested, and total RNA was prepared and analysed (fig. 4).

20 The mutants nos. 50-60 (fig. 3) show that progressive deletions of the carboxy-terminal part or the amino-terminal part of the Bt ICP coding sequence result in a gradually increasing neo transcript level. As there are not very abrupt changes in transcript
25 levels, these results suggest that the low transcript level produced by full length Bt ICP genes is controlled by a number of signals. Deletions within the Bt ICP coding sequence indeed did not localise a specific sequence element which, by itself, is
30 responsible for the low Bt ICP mRNA level. Similarly, cloning of fragments of the Bt ICP coding sequence in pJD50 (fig. 3) did not allow identification of such a region.

The relative transcript levels were plotted against the length of the Bt ICP sequence present in

the different deletion derivatives. Fig. 5 suggests that hybrid Bt ICP-ne transcript levels drop with increasing length of the Bt ICP sequence. In this respect, the mutants nos. 61-66 (fig. 3) form an exception as they show in average a low transcript level relative to the length of the Bt ICP sequence.

These results show that the low transcript levels of Bt ICP plant genes in tobacco are not exclusively due to an impaired elongation of the nascent transcript but that a number of signals operate to cause a reduced expression capacity of the chimeric Bt ICP genes.

Example 4.

To determine whether cytoplasmic events are important in causing inefficient expression of the bt2 gene in plants, the following test was carried out. Cytoplasmic bt2 mRNA steady state levels in transgenic leaf protoplasts of N28-220 are normally found to be below 1 transcript per cell. The steady state level is determined by, and is proportional to, the number of bt2 transcripts entering per time unit the cytoplasm and the cytoplasmic half-life of the transcript. When steady state levels are achieved, the absolute numbers of transcripts entering and leaving the cytoplasmic bt2 mRNA pool are equal. Therefore, the cytoplasmic half-life and cytoplasmic steady state level of the bt2 transcript will reveal whether its cytoplasmic steady state level is due to a relatively low import of bt2 transcript, a relatively high turnover (i.e., conversion to a protein) rate, or a combination of both.

The cytoplasmic turnover of bt884 transcripts was determined according to Gallie et al. (1989). A capped and polyadenylated synthetic bt884 mRNA was produced in vitro according to protocols of Promega Corporation

(Madison, Wisconsin, USA) and introduced into tobacco leaf protoplasts simultaneously with a synthetic bar (De Block et al., 1987) mRNA. The two synthetic transcripts differed only in their coding sequences. At various times after RNA delivery, samples were taken, and total RNA was isolated. Northern analyses revealed that the half-lives ($T_{1/2}$) of the synthetic bt884 and bar transcripts were about 8 ± 3 hours and 5 ± 2 hours, respectively. See Table 1, below. These data show that the bt884 coding sequence, more particularly the bt884 codon usage and the AU-rich motifs in the bt884 coding sequence, do not render the bt884 mRNA more unstable than the bar mRNA which is known to accumulate in the cytoplasm to about 1000 transcripts per tobacco leaf protoplast (calculated from Cornelissen, 1989). The low cytoplasmic steady state level of the bt884 transcripts is, therefore, caused by a lack of import of transcripts into the cytoplasm. Thus, the expression defect of the bt884 gene has to be restored by introduction of modifications in the bt884 coding sequence that improve the expression pathway in the nucleus.

Expression of the bt14, bt15 and bt18 genes in tobacco revealed that these genes also direct low cytoplasmic mRNA steady state levels. Therefore, a similar analysis was carried out with synthetic bt14, bt15 and bt18 transcripts to identify whether the expression defect had a cytoplasmic or nuclear character. Table 1, below, shows that all three transcripts behave as stable mRNAs in the cytoplasm of tobacco leaf protoplasts. Therefore, bt14, bt15 and bt18 genes, like the bt884 gene, must be deficient in exporting high levels of bt transcript to the cytoplasm, and to improve the expression of such genes, it is necessary to modify their coding sequences so

that nuclear events, which interfere with efficient gene expression, are avoided or ameliorated.

Table 1

5

Half-life determination of synthetic bt and bar mRNAs in Nicotiana tabacum cv. Petite Havanna SR1 leaf protoplasts

10	Example	1 st mRNA	T1/2 (Hours)	2 nd mRNA	T1/2 (Hours)
	A	<u>bt884</u>	8+/-3	<u>bar</u>	5+/-2
	B	<u>bt14</u>	7+/-2	<u>bar</u>	6+/-3
	C	<u>bt15</u>	12+/-5	<u>bar</u>	21+/-12
15	D	<u>bt18</u>	10+/-5	<u>bar</u>	12+/-5

Legend

20 The synthetic bar transcripts had a length of 783 bases and included a cap, the TMV leader (77 bases, Danthinne and Van Emmelo, 1990), the bar coding sequence (552 bases; De Block et al., 1987), a trailer of 52 nucleotides consisting of the bases GAUCA CGCGA AUU and 39 bases from the pGEM-3Z (Promega) polylinker (KpnI (T4 DNA pol.)-HindIII (T4 DNA pol.), and a poly(A) of 25 the composition (A)₃₃G(A)₃₂G(A)₃₂, followed by the nucleotides GCU.

30 The synthetic bt884 transcripts had a length of 2066 bases and included a cap, the TMV leader (77 bases), the bt884 coding sequence followed by the trailer until the Klenow treated PstI site (1843 nucleotides), the trailer continued with AAUUC CGGGG AUCAA UU, 39 bases of the pGEM-3Z polylinker and the (A)₃₃G(A)₃₂G(A)₂₁ poly(A), followed by the nucleotides CG.

The synthetic bt14 transcripts had a length of 2289 bases and included a cap, the TMV leader (77 bases), the bt14 coding sequence till the Klenow treated BclI site (2023 bases), plus 26 supplementary nucleotides CG UCG ACC UGC AGC CAA GCU UGC UGA, a trailer starting with UUGAU UGACC GGAUC CGGCU CUAGA AUU, followed by 39 bases of the pGEM-3Z polylinker, and the (A)₃₃G(A)₃₂G(A)₂₁ poly(A), followed by the nucleotides CGGUA CCC.

10

The synthetic bt15 transcripts had a length of 2198 bases and included a cap, the TMV leader (77 bases) the bt15 coding sequence as in pVE35 (PCT publication WO90/15139) followed by the trailer till the Klenow treated BamHI site (1989 bases), the trailer then continued with AAUU, 39 bases of the pGEM-3Z polylinker and the (A)₃₃G(A)₃₂G(A)₂₁ poly(A), followed by the nucleotides CG.

15

The synthetic bt18 transcripts had a length of 2184 bases and included a cap, the TMV leader (77 bases) the bt18 coding sequence until the Klenow treated BclI site (1918 bases), followed by 26 nucleotides until the translation stop CG UCG ACC UGC AGC CAA GCU UGC UGA, a trailer starting with UUGAU UGACC GGAUC GAUCC GGCUC AGAUC AAUU, 39 bases of the pGEM-3Z polylinker and the (A)₃₃G(A)₃₂G(A)₂₁ poly(A), followed by the nucleotides CG.

25

Example 5. Construction of Modified Bt ICP Genes

30

Examples 1-4 show that the expression in a plant of a Bt ICP gene is negatively affected by the Bt ICP coding sequence at both transcriptional and post-transcriptional levels, but principally by nuclear events. These examples also show that the control of expression is not confined to a specific DNA sequence

within the Bt ICP coding sequence. In addition, the negative effect on gene expression is an intrinsic property of the Bt ICP coding sequence. On this basis, it is believed that, by directed change of the DNA sequence of the Bt ICP coding region, an improvement of gene expression will occur. The improvement will be of a cumulative type as the negative influence of the Bt ICP coding region is spread over the complete coding sequence. Similarly, an improvement of gene expression will be obtained by reduction of the length of the Bt ICP coding sequence. This improvement will have a cumulative effect if used in combination with modifications of the Bt ICP coding region.

Therefore, two types of modifications were introduced into a Bt ICP (i.e., bt2) coding sequence which, as will be shown, indeed resulted in a significant increase in Bt ICP plant gene expression. First, the DNA sequence was modified in the central region of the toxic core fragment of the Bt ICP as transcription elongation is impaired in this region. Secondly, the length of the Bt ICP coding sequence was reduced as the negative influence is proportional to the length of the Bt ICP coding sequence. A detailed description of the mutations is given in figs. 6a, b and c. As shown in fig. 7, the modifications change the AT-content of the chimaeric Bt ICP gene significantly. The modifications change the primary DNA structure of the Bt ICP coding sequence without affecting the amino acid sequence of the encoded protein. It is evident that, if more DNA mutations were to be introduced into the Bt ICP coding sequence, a further improvement of gene expression would be obtained.

To determine the effect of the modifications, the expression properties of the modified BtICP gene and the parental bt860-neo gene were compared in a

transient expression system as described by Corn lissen and Vandewiele (1989) and Denecke et al. (1989). Basically, the accumulation profiles of the genes under study were compared by relating their profiles to the profile of a reference gene present in the same experiment. Fig. 8a shows the vectors used in the assay, and fig. 8b shows that the accumulation of the reference CAT protein is nearly identical in both experiments. It is not possible to measure the accumulation of Bt ICP encoded by the parental ~~bt860-neo~~ gene, but the modified Bt ICP gene clearly directs an increased synthesis of Bt ICP.

These results demonstrate that mutation of the Bt ICP coding sequence relieves the negative influence of the Bt ICP coding sequence on the expression of a Bt ICP plant gene.

Example 6. Cloning and Expression of Modified BT ICP Genes in Tobacco and Potato Plants

Using the procedures described in US patent application 821,582, filed January 22, 1986, and EPA 86300291.1, EPA 88402115.5 and EPA 89400428.2, the modified Bt ICP (i.e., bt2) genes of figs. 6 and 7 are inserted into the intermediate T-DNA vector, pGSH1160 (Deblaere et al., 1988) between the vector's T-DNA terminal border repeat sequences.

To obtain significant expression in plants, the modified Bt ICP genes are placed under the control of the strong TR2' promoter (Velten et al., 1984) and are fused to the transcript 3' end formation and polyadenylation signals of the T-DNA gene 7 (Velten and Schell, 1985).

In addition, the translation initiation context is changed in accordance with the Japanese consensus sequence (Joshi, 1987) in order to optimize the

translati n initiati n in plant c lls. To this end, an olig duplex (figs. 6a and 6b) is introduced t cr at the following sequence at translation initiation site: AAAACCATGGCT. In this way, an additional codon (i.e.,
5 GCT) coding for alanine is introduced. Additionally, KpnI and BstXI sites are created upstream of the ATG translation initiation codon.

Using standard procedures (Deblaere et al., 1985), the intermediate plant expression vectors, containing
10 the modified BtICP gene, are transferred into the Agrobacterium strain C58C1 D1⁸ (US patent application 821,582; EPA 86300291.1) carrying the disarmed Ti-plasmid pGV2260 (Vaeck et al., 1987). Selection for spectinomycin resistance yields cointegrated plasmids,
15 consisting of pGV2260 and the respective intermediat plant expression vectors. Each of these recombinant Agrobacterium strains is then used to transform different tobacco plant cells (Nicotiania tabacum) and potato plant cells (Solanum tuberosum) so that the
20 modified Bt ICP genes are contained in, and expressed by, different tobacco and potato plant cells.

The transgenic tobacco plants containing the modified Bt ICP genes are analyzed with an ELISA assay. These plants are characterized by a significant
25 increase in levels of Bt (Bt2) proteins, compared to a transgenic tobacco plant containing a non-modified Bt ICP (bt2) gene.

The insecticidal activity of the expression products of the modified Bt ICP (bt2) genes in leaves
30 of transformed tobacco and potato plants is evaluated by recording the growth rate and mortality of larvae of Tobacco hornworm (Manduca sexta), Tobacco budworm (Heliothis vir scens) and potat tubermoth (Phthorimaea operculella) fed on leaves of these tw types of plants. Thes results are compared with the gr wth rat

of larva fed leaves from tobacco and potato plants transformed with the unmodified or parental Bt ICP (bt2) gene and from untransformed potato and tobacco plants. Toxicity assays are performed as described in EPA 88402115.5 and EPA 86300291.1.

A significantly higher mortality rate is obtained among larvae fed on leaves of transformed plants containing and expressing the modified Bt ICP genes. Tobacco and potato plants containing the modified Bt ICP genes show considerably higher expression levels of Bt ICPs compared to tobacco and potato plants containing the unmodified Bt ICP gene.

The insecticidal activity of three transgenic tobacco plants containing the modified Bt ICP genes is determined against second and third instar larvae of Heliothis virescens. The control plant was not transformed. The results are summarized in Table 2, below.

Table 2

Plant	% mortality of insects (recorded after 5 days)
Control	11
No. 1	100
No. 2	88.5
No. 3	100

Needless to say, this invention is not limited to tobacco and potato plants transformed with the modified Bt ICP gene. It includes any plant, such as tomato, alfalfa, sunflowers, corn, cotton, soybean, sugar beets, rapeseed, brassicas and other vegetables, transformed with the modified Bt ICP gene.

Nor is the invention limited to the use of Agrobacterium tumefaciens Ti-plasmids for transforming plant cells with a modified Bt ICP gene. Other known techniques for plant transformation, such as by means of liposomes, by electroporation or by vector systems based on plant viruses or pollen, can be used for transforming monocotyledonous and dicotyledonous with such a modified Bt ICP gene.

Nor is the invention limited to the bt2 gene, but rather encompasses all Cry I, Cry II, CryIII and Cry IV Bt ICP genes.

15

20

25

30

References

- Adang et al., Gene 36, 289-300 (1985).
- Asselin et al., Oncogene 4, 549-558 (1989).
- 5 - Barton et al., Plant Physiol. 85, 1103-1109 (1987).
- Connely and Manley, Cell 57, 567-571 (1989)
- Connely and Manley, Molecular and Cellular Biology, 9, 5254-5259 (1989)
- Cornelissen, Nucl. Acids Res. 17, 7203-7209 (1989).
- 10 - Cornelissen and Vandewiele, Nucl. Acids Res. 17,
2323-2343 (1989).
- Cox and Goldberg, in: Plant Molecular Biology, A practical approach, edited by C. Shaw, IRL Press, pp.1-35 (1988).
- 15 - Danthinne and Van Emmelo, Me. Fac. Landbouww. Rijksuniv. Gent 55 (3a), 1037-1045 (1990).
- Darnell, Nature 297, 365-371 (1982).
- Deblaere et al., Methods in Enzymology 153, 277-292 (1988).
- 20 - Deblaere et al., Nucleic Acids Research 13, 4777-4787 (1985).
- De Block et al., EMBO J. 6, 2513-2518 (1987).
- Delannay et al., Bio/Technology 7, 1265-1269 (1989).
- Denecke et al., Meth. Mol. Cel. Biol. 1, 19-27 (1989).
- 25 - Diffley and Stillman, Mol. and Cell. Biol. 6, 1363-1373 (1986).
- Dulmage H. T, "Production of Bacteria for Biological Control of Insects" in "Biological Control in Crop Production" Ed. Paparizas D. C., Osmum Publishers, Totowa, N.Y., USA, pp. 129 - 141 (1981).
- 30 - Engler et al., J. Biol. Chem. 263, 12384-12390 (1988).
- Engvall and Pesc , Scand. Immun 1. Suppl. 7 (1978).
- Fischhoff et al., Bi /Techno logy 5, 807-812 (1987).

- Fromm et al., Bio/Technology 8, 833 et seq. (1990).
- Galli et al., Plant Cell 1, 301-311 (1989).
- Gasser and Fraley, Science 244, 1293-1299 (1989).
- Gielen et al., EMBP J 3, 835-845 (1984).
- 5 - Goldberg L. and Margalit J., Mosq. News 37, 355-358 (1977).
- Gordon-Kamm et al., The Plant Cell 2, 603 et seq. (1990).
- Gould et al., Plant Physiol. 95, 426-434 (1991).
- 10 - Hoffmann et al., Proc. Natl. Acad. Sci. USA 85,
7844-7848 (1988).
- Höfte et al., Eur. J. Biochem 161, 273-280 (1986).
- Höfte H. and Whiteley H. R., Microbiological Reviews 53, 242-255 (1989).
- 15 - Joshi, (1987) Nucl. Acids Res. 15, 6643-6653.
- Krieg et al., Z. Ang. Ent 96, 500-508 (1983).
- Luthe and Quatrano, Plant and Physiology 65, 305 (1980).
- Mc Pherson et al., Bio/Technology 6, 61-66 (1988).
- 20 - Meeusen and Warren, Ann. Rev. Entomol. 34, 373-381 (1989).
- Miller et al., Molecular and Cellular Biology 9, 5340-5349 (1989).
- "The Molecular Biology of the Cell, Second Edition",
25 Eds. Alberts et al., Garland Publishing Co., N.Y. and London (1989).
- Neumann et al., Biotechniques 5, 144 (1987).
- Odell et al., Nature 313, 810-812 (1985).
- Schnepf et al., J. Biol. Chem. 260, 6264-6272 (1985).
- 30 - Vaeck et al., Nature 327, 33-37 (1987).
- Velten and Schell, Nucleic Acids Research 13, 6981-6998 (1985).
- Velten et al., EMBO J 3, 2723-2730 (1984).
- Widner and Whiteley, J. Bacterial 171, 965-974 (1989).

43

- Yanisch-Perron et al., Gene 33, 103-109 (1985).

5

10

15

20

25

30

Claims

1. A process for modifying a Bt ICP gene to improve its expression in a plant cell, transformed with the gene; the process comprising the step of: changing A and T sequences in a plurality of translational codons of the gene to corresponding G and C sequences encoding the same amino acids, so as to improve the gene's transcription to an mRNA, the nuclear accumulation of the mRNA and/or the nuclear export of the mRNA, particularly the gene's transcription, in the plant cell.
2. The process of claim 1 for modifying a Bt ICP gene to improve its transcription in plant cells, transformed with the modified gene, wherein the plurality of translational codons is at least one region of the gene which, during transcription, has thereon a relatively low percentage of RNA polymerase II of the plant cell as compared to another adjacent upstream region of the gene.
3. The process of claim 1 or 2, wherein the Bt ICP gene encodes a Bt insecticidal crystal protein truncated towards a trypsin cleavage site, preferably at both its C-terminal and N-terminal ends, and preferably encoding a portion of the protein of about 60 - 80 kDa, particularly the toxin of the protein.
4. The process of anyone of claims 1-3, wherein A and T sequences of at least 3 codons are changed to G and C sequences at a translation initiation site of the gene and A and T sequences of at least about 3, preferably at least about 10, especially at least about 33, codons are changed to G and C sequences in a second region of the gene, preferably affecting transcription

longati n, downstr am of th translation initiati n site.

5. The process of anyone of claims 1-4, wherein A and
5 T sequences of at least about 3, preferably at least about 10, especially at least about 33, codons are changed to G and C sequences in a third region of the gene, preferably affecting cytoplasmic RNA concentration.
- 10 6. The process of claim 4 or 5, wherein A and T sequences of at least about 3 codons are changed to G and C sequences at a translation termination end of the gene.
- 15 7. The process of anyone of claims 4-6, wherein the gene is a cryI gene, such as a bt2, bt14, bt15 or bt18 gene, preferably a bt2 gene, or a gene having substantial sequence homology thereto.
- 20 8. The process of claim 7 wherein the gene is a bt2 gene; the second region being between about nucleotides 674 and 1000 and A and T sequences of about 59 or more codons are changed to G and C sequences in the second region, particularly between about nucleotides 733 and
25 1000, quite particularly between about nucleotides 765 and 794.
9. The process of anyone of claims 1-8, wherein the gene is further modified by substituting for its ATG
30 translation initiation site : AAAACCATGGCT.
10. The modified Bt ICP gene obtained by the process of anyone of claims 1-9.

11. A chimaeric gene for transforming a cell of a plant, comprising the following operably-linked DNA fragments in the same transcriptional unit:

- 5 a) the modified Bt ICP gene of claim 10;
- b) a promoter capable of directing expression of the modified Bt ICP gene in the plant cell; and
- 10 c) transcript 3' end formation and polyadenylation signals suitable for expressing the modified Bt ICP gene in the plant cell.

12. The plant cell of claim 11, transformed with the chimaeric gene of claim 11.

- 15 13. A plant, plant tissue or plant cell culture consisting of the plant cells of claim 12.

14. A seed of the plant of claim 13.

- 20 15. A vector, preferably a Ti-plasmid, for stably transforming the nuclear genome of a plant, comprising the chimaeric gene of claim 11.

- 25 16. A process for protecting the plant of claim 10 against an insect pest, comprising the step of: transforming the genome of the plant with the chimaeric gene of claim 11.

- 30 17. A process for modifying a foreign gene whose rate and/or level of expression in a plant cell, transformed with the gene, is substantially limited by the rate and/or level of nuclear production of an mRNA encoded by the gene; the process comprising the step of: changing A and T sequences in a plurality of translational codons of the gene, particularly in a plurality of translational codons in at least one

region of the gene which, during transcription, has thereon a relatively low percentage of RNA polymerase II of the plant cell as compared to another adjacent upstream region of the gene; the A and T sequences
5 being changed to corresponding G and C sequences encoding the same amino acids, so as to improve the gene's transcription to the mRNA, the nuclear accumulation of the mRNA and/or the nuclear export of the mRNA, particularly the gene's transcription to the
10 mRNA, quite particularly the transcript elongation by RNA polymerase II on the gene, in the plant cell.

15

20


25

30

Figure 1

ID PGSH163 : pgsh163

Showing 4323 out of 10457 bases; 41% of the molecule is visible.



3/26

FIGURE 2a (cont)

Restriction fragment	size (bp)	# UTP	fraction #UTP size (bp)	SCANNING VALUE AUTO-RADIOGRAM	X
BamHI-HindIII	1700	513	0.302	6565	12.8
BamHI-SacI	1353	412	0.305	4572	11.0
BamHI-EcoRV	730	229	0.314	2466	10.6
BamHI-EcoRI	275	96	0.349	-	-
XmnI-XmnI	628	173	0.275	1817	10.5
EcoRI-EcoRI	725	224	0.309	2926	13.1
XmnI-XmnI	729	212	0.291	736	3.47
EcoRV-EcoRV	840	249	0.296	1102	4.43
EcoRI-EcoRI	585	149	0.255	635	4.26

4/26






FIGURE 2b

Figure 2cRNA polymerase II migration over
the bt2 coding sequence

6/26

FIGURE 3

SPLICED PVE36, SEQ 1:8600 INTO PVE36, SEQ
8600 BASE PAIRS




1 : AatII	8 : BclI	15 : EcoRV	22 : NarI	29 : RsrII
2 : AccI	9 : BspMII	16 : Eco31I	23 : NdeI	30 : SacI
3 : AflII	10 : BssHII	17 : EspI	24 : NheI	31 : SpeI
4 : AlwI	11 : BstEII	18 : HindII	25 : NsiI	32 : SphI
5 : ApaI	12 : BstXI	19 : HindIII	26 : PflMI	33 : Tth111I
6 : AvrII	13 : EagI	20 : KpnI	27 : PpuMI	34 : XcaI
7 : BamHI	14 : EcoNI	21 : NaeI	28 : PvuI	35 : XhoI

SUBSTITUTE SHEET

7/26


FIGURE 3(CONT.)



SUBSTITUTE SHEET

9/26

FIGURE 3(CONT.)



10/26


sample	gene assayed	ug loaded neo/cat	cpm (neo)	pg (cat)	cpm (cat)	pg (cat)	neo- SR1	neo/ cat	% rel to pJD51
pJD50	nptII/cat	4/1	121	1.366	32	0.283	1.279	4.52	16.53
pJD51	nptII/cat	4/1	131.3	1.482	20.7	0.051	1.395	27.35	100
pJD52	nptII/cat	4/1	50.3	0.568	23.3	0.104	0.84	4.62	16.89
pJD53	nptII/cat	4/1	44.3	0.5	36.7	0.379	0.431	1.09	3.99
pJD54	nptII/cat	4/1	22	0.248	28.3	0.207	0.161	0.78	2.85
pJD55	nptII/cat	4/1	374.3	4.225	35	0.344	4.138	12.03	43.99
pJD56	nptII/cat	4/1	13.3	0.15	38	0.406	0.063	0.16	0.59
pJD57	nptII/cat	4/1	55.7	0.629	21.7	0.071	0.542	7.63	27.9
pJD58	nptII/cat	4/1	78.3	0.883	29.7	0.236	0.797	3.38	12.36
pJD59	nptII/cat	4/1	17.3	0.195	38.3	0.412	0.108	0.26	0.95
pJD60	nptII/cat	4/1	15.7	0.177	29	0.221	0.09	0.41	1.5
pJD61	nptII/cat	4/1	286	3.228	42.3	0.494	3.141	6.36	23.25
pJD62	nptII/cat	4/1	186.7	2.108	69	1.042	2.021	1.94	7.09
pJD63	nptII/cat	4/1	117.7	1.329	43.3	0.515	1.242	2.41	8.81
pJD64	nptII/cat	4/1	188.7	2.13	57.7	0.81	2.043	2.52	9.21
pJD65	nptII/cat	4/1	158.7	1.791	42	0.488	1.704	3.49	12.76
pJD66	nptII/cat	4/1	72.3	0.816	65	0.96	0.73	0.76	2.78
pPS56D1	nptII/cat	4/1	24	0.271	56.7	0.79	0.184	0.23	0.84
pPS56D2	nptII/cat	4/1	59.3	0.669	100	1.68	0.583	0.35	1.28
pPS56D3	nptII/cat	4/1	214.3	2.419	49.3	0.64	2.332	3.66	13.38
pPS56D4	nptII/cat	4/1	54.3	0.613	29.7	0.236	0.526	2.23	8.15
pPS56D6	nptII/cat	4/1	37	0.418	46.7	0.584	0.331	0.57	2.08
pPS56D7	nptII/cat	4/1	35.3	0.398	31.3	0.268	0.311	1.16	4.24
pPS56D8	nptII/cat	4/1	43.3	0.489	45.7	0.564	0.402	0.71	2.6

Figure 4

11/26

Effect of the length of bt2 sequences on bt2:neo derived mRNA

Figure 5




OLIGOPS15

GTACCAAACCATGGCTATCGAGACCGGTTACACCCCAATCGATATCG

OLIGOPS16

ATCGATTGGGGTGTAACCGGTCTCGATAGCCATGGTTTTGGTACCGAT

Figure 6a(cont.)

3) OLIGOPS16', 4) OLIGOPS15,

Name	Base	
3	1	ATCGGTACCA AAACCATGGC TATCGAGACC GGTTACACCC CAATCGAT
4	1	GTACCA AAACCATGGC TATCGAGACC GGTTACACCC CAATCGATAT CG
CON	1	ATCGGTACCA AAACCATGGC TATCGAGACC GGTTACACCC CAATCGATAT CG

14/26

Figure 6a(cont.)

ATC GGT ACC AAA ACC ATG GCT ATC GAG ACC GGT TAC ACC CCA ATC GAT ATC G
MET Ala Ile Glu Thr Gly Tyr Thr Pro Ile Asp Ile

OLIG01

GATCCTCTAGAGACTGGATCAGGTACAACCAAGTTCAGGAGGGAGTTAACCCCTAACCGTGTTAGAC

OLIG02

ATCGTGTCCCTATTCCCGAACTACGACAGCAGGACGTACCCAATCCGAACCGTGTTCCAGTTAACAGGGA

OLIG03

GATCTACACCAACCCAGTGTTAGAGAACTTCGACGGTAGCTTCCGAGGCTCGGCTCAGGGCATCG

OLIG04

AGGGAAGCATCAGGAGCCACACTTGATGGACATCCTTAACAGCATCACCATCTACACGGACGCT

OLIG05

CACAGGGGAGAGTACTACTGGTCCGGGCACCAGATCATGGCTTCCCCTGTGGGGTTCTCGGGGCCAGAATTCC

OLIG06

GATCCGAATTCTGGCCCCGAGAACCCACAGGGGAAGCCATGATCTGGTGCCCGGACCAGTAGTAC

OLIG07

TCTCCCCTGTGAGCGTCCGTGTAGATGGTGATGCTGTTAAGGATGTCCATCAAGTGTGGGCTCCT

OLIG08

GATGCTTCCCTCGATGCCCTGAGCCGAGCCTCGGAAGCTACCGTCGAAGTTCTCTAACACTGGG

OLIG09

TTGGTGTAGATCTCCCTGGTTAACTGGGACACGGTTCGGATTGGGTACGTCCCTGCTGTCGTAGTTCGGGAA

OLIG010

TAGGGACACGATGTCTAACACGGTTAGGGTTAACTCCCTCCTGAACTGGTTGTACCTGATCCAGTCTCTAGAC

SUBSTITUTE SHEET

synth -9

Linear

LENGTH - 343

			149	213	278	
	77		148	212	277	
5	66		137	202	267	34
1	65 78		136	201	266	33

#-----#
 <--OLIGO10'---<<--OLIGO9'---<--OLIGO8'---<<--OLIGO7'---<--OLIGO6'---<
 >--OLIGO1--->--OLIGO2--->--OLIGO3--->--OLIGO4--->--OLIGO5--->

Name Base

1	1	GATCCTCTAG	AGACTGGATC	AGGTACAACC	AGTTCAGGAG	GGAGTTAACC	CTAACCGTGT
10	1	CTCTAG	AGACTGGATC	AGGTACAACC	AGTTCAGGAG	GGAGTTAACC	CTAACCGTGT

CON	1	GATCCTCTAG	AGACTGGATC	AGGTACAACC	AGTTCAGGAG	GGAGTTAACC	CTAACCGTGT
-----	---	------------	------------	------------	------------	------------	------------

1	61	TAGAC					
10	57	TAGACATCGT	GTCCCTA				
2	1	ATCGT	GTCCCTATTC	CCGAACTACG	ACAGCAGGAC	GTACCCAATC	CGAACCGTGT
9	1		TTC	CCGAACTACG	ACAGCAGGAC	GTACCCAATC	CGAACCGTGT

CON	61	TAGACATCGT	GTCCCTATTC	CCGAACTACG	ACAGCAGGAC	GTACCCAATC	CGAACCGTGT
-----	----	------------	------------	------------	------------	------------	------------

2	56	CCCAGTTAAC	CAGGGA				
9	44	CCCAGTTAAC	CAGGGAGATC	TACACCAA			
3	1		GATC	TACACCAACC	CAGTGTTAGA	GAAGTTTCGAC	GGTAGCTTCC
8	1			CC	CAGTGTTAGA	GAAGTTTCGAC	GGTAGCTTCC

CON	121	CCCAGTTAAC	CAGGGAGATC	TACACCAACC	CAGTGTTAGA	GAAGTTTCGAC	GGTAGCTTCC
-----	-----	------------	------------	------------	------------	-------------	------------

3	45	GAGGCTCGGC	TCAGGGCATC	G			
8	33	GAGGCTCGGC	TCAGGGCATC	GAGGGAAGCA	TC		
4	1			AGGGAAGCA	TCAGGAGCCC	ACACTTGATG	GACATCCTTA
7	1				AGGAGCCC	ACACTTGATG	GACATCCTTA

CON	181	GAGGCTCGGC	TCAGGGCATC	GAGGGAAGCA	TCAGGAGCCC	ACACTTGATG	GACATCCTTA
-----	-----	------------	------------	------------	------------	------------	------------

4	40	ACAGCATCAC	CATCTACACG	GACGCT			
7	29	ACAGCATCAC	CATCTACACG	GACGCTCACA	GGGGAGA		
5	1			CACA	GGGGAGAGTA	CTACTGGTCC	GGGCACCAGA
6	1				GTA	CTACTGGTCC	GGGCACCAGA

CON	241	ACAGCATCAC	CATCTACACG	GACGCTCACA	GGGGAGAGTA	CTACTGGTCC	GGGCACCAGA
-----	-----	------------	------------	------------	------------	------------	------------

5	35	TCATGGCTTC	CCCTGTGGGG	TTCTCGGGGC	CAGAATTCG		
6	24	TCATGGCTTC	CCCTGTGGGG	TTCTCGGGGC	CAGAATTCGG	ATC	

CON	301	TCATGGCTTC	CCCTGTGGGG	TTCTCGGGGC	CAGAATTCGG	ATC	
-----	-----	------------	------------	------------	------------	-----	--

SYNTHE-9

27

54

GAT CCT CTA GAG ACT GGA TCA GGT ACA ACC AGT TCA GGA GGG AGT TAA CCC TAA
 Ser S r Arg Asp Trp Ile Arg Tyr Asn Gln Phe Arg Arg Glu Leu Thr Leu Thr

81

108

CCG TGT TAG ACA TCG TGT CCC TAT TCC CGA ACT ACG ACA GCA GGA CGT ACC CAA
 Val Leu Asp Ile Val Ser Leu Phe Pro Asn Tyr Asp Ser Arg Thr Tyr Pro Il

135

162

TCC GAA CCG TGT CCC AGT TAA CCA GGG AGA TCT ACA CCA ACC CAG TGT TAG AGA
 Arg Thr Val Ser Gln Leu Thr Arg Glu Ile Tyr Thr Asn Pro Val Leu Glu Asn

189

216

ACT TCG ACG GTA GCT TCC GAG GCT CGG CTC AGG GCA TCG AGG GAA GCA TCA GGA
 Phe Asp Gly Ser Phe Arg Gly Ser Ala Gln Gly Ile Gln Gly Ser Ile Arg Ser

243

270

GCC CAC ACT TGA TGG ACA TCC TTA ACA GCA TCA CCA TCT ACA CGG ACG CTC ACA
 Pro His Leu MET Asp Ile Leu Asn Ser Ile Thr Ile Tyr Thr Asp Ala His Arg

297

324

GGG GAG AGT ACT ACT GGT CCG GGC ACC AGA TCA TGG CTT CCC CTG TGG GGT TCT
 Gly Glu Tyr Tyr Trp Ser Gly His Gln Ile MET Ala Ser Pro Val Gly Phe S r

CGG GGC CAG AAT TCG GAT C
 Gly Pro Glu Phe Gly

Figure 6c

PVE36.SEQ nucleotide 800 - 4000

AAATGGATAAATAGCCTTGCTTCCTATTATATCTTCCCAAATTACCAATACATTACACTAGCATCTGAAT
TTCATAACCAATCTCGATACACCAAATCGATGGATCCCGATAACAATCCGAACATCAATGAATGCATTCC
TTATAATTGTTTAAGTAACCCCTGAAGTAGAAGTATTAGGTGGAGAAAGAATAGAAACTGGTTACACCCCA
ATCGATATTTCTTGTCGCTAACGCAATTTCTTTTGAGTGAATTTGTTCCCGGTGCTGGATTTGTGTTAG
GACTAGTTGATATAATATGGGGAATTTTGGTCCCTCTCAATGGGACGCATTTCTTGACAAATTGAACA
GTTAATTAACCAAGAATAGAAGAATTCGCTAGGAACCAAGCCATTTCTAGATTAGAAGGACTAAGCAAT
CTTTATCAAATTTACGCAGAATCTTTTAGAGAGTGGGAAGCAGATCCTACTAATCCAGCATTAAGAGAAG
AGATGCGTATTCAATTCAATGACATGAACAGTGCCTTACAACCGCTATTCTCTTTTGCAGTTCAAAA
TTATCAAGTTCCTCTTTTATCAGTATATGTTCAAGCTGCAATTTACATTTTATCAGTTTTGAGAGATGTT
TCAGTGTGTTGGACAAAGGTGGGGATTTGATGCCGCGACTATCAATAGTCGTTATAATGATTTAACTAGGC
TTATTGGCAACTATACAGATCATGCTGTACGCTGGTACAATACGGGATTAGAGCGTGTATGGGGACCGGA
TTCTAGAGATTGGATAAGATATAATCAATTTAGAAGAGAATTAACACTAACTGTATTAGATATCGTTTCT
CTATTTCCGAACATGATAGTAGAAGCTATCCAATTCGAACAGTTTCCCAATTAACAAGAGAAATTTATA
CAAAACCCAGTATTAGAAAATTTTATGGTAGTTTCGAGGCTCGGCTCAGGGCATAGAAGGAAGTATTAG
GAGTCCAGATTTGATGGATATACTTAACAGTATAACCATCTATACGGATGCTCATAGAGGAGAATATTAT
TGGTCAGGGCATCAAATAATGGCTTCTCCTGTAGGGTTTTCGGGGCCAGAATTCATTTTTCCGCTATATG
GAACATATGGGAAATGCAGCTCCACAACAACGATTTGTTGCTCAACTAGGTCAGGGCGTGTATAGAACATT
ATCGTCCACTTATATAGAAGACCTTTAATATAGGGATAAATAATCAACAACATATCTGTTCTTGACGGG
ACAGAATTTGCTTATGGAACCTCCTCAAATTTGCCATCCGCTGTATACAGAAAAAGCGGAACGGTAGATT
CGCTGGATGAAATACCGCCACAGAATAACAACGTGCCACCTAGGCAAGGATTTAGTCATCGATTAAAGCCA
TGTTTCAATGTTTCGTTCAGGCTTTAGTAATAGTAGTGAAGTATAATAAGAGCTCCTATGTTCTCTTGG
ATACATCGTAGTGCTGAATTTAATAATATAATTCCTTCATCACAATTAACAAATACCTTTAACAAAAT
CTACTAATCTTGGCTCTGGAACCTCTGTGCTTAAAGGACAGGATTTACAGGAGGAGATATTCTTCGAAG
AACTTCACCTGGCCAGATTTCAACCTTAAGAGTAAATATTACTGCACCATTATCACAAAGATATCGGGTA
AGAATTCGCTACGCTTCTACCACAAATTTACAATTCATACATCAATTGACGGAAGACCTATTAAATCAGG
GGAATTTTTTACGAACTATGAGTAGTGGGAGTAATTTACAGTCCGGAAGCTTTAGGACTGTAGGTTTTAC
TACTCCGTTTAACTTTTCAAATGGATCAAGTGTATTTACGTTAAGTGCTCATGTCTTCAATTCAGGCAAT
GAAGTTTATATAGATCGAATTGAATTTGTTCCGGCAGAAGTAACCTTTGAGGCAGAATATGATTTAGAAA
GAGCACAAAAGGCGGTGAATGAGCTGTTTACTTCTTCCAATCAAATCGGGTTAAAAACAGATGTGACGGA
TTATCATATTGATCAAGTATCCAATTTAGTTGAGTGTATCTGATGAATTTTGTCTGGATGAAAAAAA
GAATTGTCCGAGAAAGTCAAACATGCGAAGCGACTTAGTGATGAGCGGAAXXXXCCTCGAGCTTGGATG
GATTGCAACGAGGTTCTCCGGCCGCTTGGGTGGAGAGGCTATTCCGGCTATGACTGGGCAACAACAGACAAT
CGGCTGCTCTGATGCCGCCGTGTTCCGGCTGTACGCGCAGGGGCGCCCGGTTCTTTTTGTCAAGACCGAC
CTGTCGGGTGCCCTGAATGAACTGCAGGACGAGGCAGCGCGGCTATCGTGGCTGGCCACGACGGGCGTTC
CTGCGCAGCTGTGCTCGACGTTGTCACTGAAGCGGGAAGGGAAGTGGCTGCTATTGGGCGAAGTGCCGGG
GCAGGATCTCCTGTCTCATCTCACCTTGCTCCTGCCGAGAAAGTATCCATCATGGCTGATGCAATGCGGCGG
CTGCATACGCTTGATCCGGCTACCTGCCATTTCGACCACCAAGCGAAACATCGCATCGAGCGAGCAGCTA
CTCGGATGGAAGCCGGTCTTGTGATCAGGATGATCTGGACGAAGAGCATCAGGGGCTCGCGCCAGCCGA
ACTGTTCCGCCAGGCTCAAGGCGCGCATGCCGACGGCGAGGATCTCGTCTGACCCATGGCGATGCCTGC
TTGCCGAATATCATGGTGGAAAATGGCCGCTTTTCTGGATTCTCGACTGTGGCCGGCTGGGTGTGGCGG
ACCGCTATCAGGACATAGCGTTGGCTACCCGTGATATTGCTGAAGAGCTTGGCGGCGAATGGGCTGACCG
CTTCTCGTGCTTTACGGTATCGCCGCTCCCGATTTCGAGCGCATCGCCTTCTATCGCCTTCTTGACGAG
TTCTTCTGACAGATCCCCGATGAGCTAAGCTAGCTATATCATCAATTTATGTATTACACATAATATCGC
ACTCAGTCTTTTATCTACGGCAATGTACCAGCTGATATAATCAGTTATTGAAATATTTCTGAATTTAAAC
TTGCATCAATAAATTTATGTTTTTGTGTTGGACTATAATACCTGACTTGTTATTTTATCAATAAATATTTA
AACTATATTTCTTTCAAGATGGGAATTAACATCTACAAATTGCCTTTTCTT

The ATG initiation codon and the TGA stop codon are underlined.

19/26

Figure 6c (cont.)

PPS029.SEQ nucl otide 800 - 4000


AAATGGATAAATAGCCTTGCTTCCTATTATATCTTCCCAAATTACCAATACATTACACTAGCATCTGAAT
 TTCATAACCAATCTCGATACACCAAATCGGTACCAAACCATGGCTATCGAGACCGGTTACACCCCAATC
 GATATTTCTTGTGCTAACGCAATTTCTTTTGAGTGAATTTGTTCCCGGTGCTGGATTGTGTTAGGAC
 TAGTTGATATAATATGGGGAATTTTGGTCCCTCTCAATGGGACGCATTTCTTGTAACAAATTGAACAGTT
 AATTAACCAAAGATAGAAGAATTCGCTAGGAACCAACCATTTCTAGATTAGAAGGACTAAGCAATCTT
 TATCAAATTTACGCAGAATCTTTTAGAGAGTGGGAAGCAGATCCTACTAATCCAGCATTAAGAGAAGAGA
 TGGCTATTCAATTCAATGACATGAACAGTGCCCTTACAACCGCTATTCTCTTTTGCAGTTCAAAATTA
 TCAAGTTCTCTTTTATCAGTATATGTTCAAGCTGCAAAATTACATTTATCAGTTTGGAGAGATGTTTCA
 GTGTTTGGACAAAGGTGGGGATTGATGCCGCGACTATCAATAGTCGTTATAATGATTTAACTAGGCTTA
 TTGGCAACTATACAGATCATGCTGTACGCTGGTACAATACGGGATTAGAGCGTGTATGGGGACCGGATTC
 TAGAGACTGGATCAGGTACAACCAAGTTCAGGAGGGAGTTAACCTAACCGTGTAGACATCGTGTCCCTA
 TTCCCGAATACGACAGCAGGACGTACCCAATCCGAACCGTGTCCAGTTAACAGGGAGATCTACACCA
 ACCCAGTGTAGAGAACTTCGACGGTAGCTTCGAGGGCTCGGCTCAGGGCATCGAGGGAAGCATCAGGAG
 CCCACACTTGATGGACATCCTTAACAGCATCACCATCTACACGGACGCTCACAGGGGAGAGTACTACTGG
 TCCGGGCACAGATCATGGCTTCCCTGTGGGGTCTCGGGGCCAGAATTCATTTTCCGCTATATGGAA
 CTATGGGAAATGCAGCTCCACAACAACGTATTGTGCTCAACTAGGTCAGGGCGTGTATAGAACATTATC
 GTCCACTTTATATAGAAGACCTTTAATATAGGGATAAATAATCAACAACATATCTGTTCTTGACGGGACA
 GAATTTGCTTATGGAACTCCTCAAATTTGCCATCCGCTGTATACAGAAAAGCGGAACGGTAGATTTCG
 TGGATGAAATACCGCCACAGAATAACAACGTGCCACCTAGGCAAGGATTTAGTCAATGATTAAGCCATGT
 TTCAATGTTTCGTTTCAGGCTTTAGTAATAGTAGTGTAAGTATAATAAGAGCTCCTATGTTCTCTTGATA
 CATCGTAGTGCTGAATTTAATAATATAATTCCTTCATCACAATTAACAAATACCTTTAACAAAATCTA
 CTAATCTTGGCTCTGGAATCTGTGCTTAAAGGACCAGGATTTACAGGAGGAGATATTCTTCGAAGAAC
 TTCACCTGGCCAGATTTCAACCTTAAGAGTAAATATTACTGCACCATTATCACAAAGATATCGGGTAAGA
 ATTCGCTACGCTTCTACCACAAATTTACAATTCATACATCAATTGACGGAAGACCTATTAATCAGGGGA
 ATTTTTCAGCAACTATGAGTAGTGGGAGTAATTTACAGTCCGGAAGCTTTAGGAGCTAGGTTTACTAC
 TCCGTTTAACTTTTCAAATGGATCAAGTGTATTTACGTTAAGTGCTCATGTCTTCAATTCAGGCAATGAA
 GTTTATATAGATCGAATTGAATTTGTTCCGGCAGAAGTAACTTTGAGGCAGAATATGATTTAGAAAGAG
 CACAAAAGGCGGTGAATGAGCTGTTTACTTCTTCCAATCAAATCGGGTTAAAAACAGATGTGACGGATTA
 TCATATTGATCAAGTATCCAATTTAGTTGAGTGTTTATCTGATGAATTTTGTCTGGATGAAAAAAGAA
 TTGTCCGAGAAAGTCAAACATGCCAAGCGACTTAGTGATGAGCGGAAXXXXCCTCGAGCTTGGATGGAT
 TGCACGCAGGTTCTCCGGCCGCTTGGGTGGAGAGGCTATTCCGCTATGACTGGGCACAAACAGCAATCGG
 CTGCTCTGATGCCGCCGTGTTCCGGCTGTCAGCGCAGGGGCGCCCGGTTCTTTTGTCAAGACCGACCTG
 TCCGGTGCCCTGAATGAACTGCAGGACGAGGACGCGCGGCTATCGTGGCTGGCCACGACGGGCGTTCCTT
 GCGCAGCTGTGCTCGACGTTGTCACTGAAGCGGGAAGGGACTGGCTGCTATTGGGCGAAGTGCCGGGGCA
 GGATCTCCTGTCTCATCTCACCTTGCTCCTGCCGAGAAAGTATCCATCATGGCTGATGCAATGCGGCGGCTG
 CATACGCTTGATCCGGCTACCTGCCCATTGACCAACCAAGCGAAACATCGCATCGAGCGAGCACGTAATC
 GGATGGAAGCCGCTCTTGTGATCAGGATGATCTGGACGAAGAGCATCAGGGGCTCGCGCCAGCCGAATC
 GTTCGCCAGGCTCAAGGCGCGCATGCCGACGCGGAGGATCTCGTCTGACCCATGGCGATGCCTGCTTG
 CCGAATATCATGGTGGAAAATGGCCGCTTTTCTGGATTCATCGACTGTGGCCGGCTGGGTGTGGCGGACC
 GCTATCAGGACATAGCGTTGGCTACCCGTGATATTGCTGAAGAGCTTGGCGGCGAATGGGCTGACCGCTT
 CCTCGTGCTTTACGGTATCGCCGCTCCCGATTTCGACGCGCATCGCCTTCTATCGCCTTCTTGACGAGTTC
 TTCTGACAGATCCCCCGATGAGCTAAGCTAGCTATATCATCAATTTATGTATTACATAATATCGCACT
 CAGTCTTTTATCTACGGCAATGTACCAGCTGATATAATCAGTTATTGAAATATTTCTGAATTTAAACTTG
 CATCAATAAATTTATGTTTTGCTTGGACTATAATACCTGACTTGTTATTTTATCAATAAATATTTAAAC
 TATATTTCTTTCAAGATGGGAATTAACATCTACAAATTGCCTTTTCTATCGACCATGTACGGGTACCGA
 GCTCGAATTCCTACGCAGCAGGTCTCATCAAGACGATCTACCCGAGTAACA

the ATG initiation codon and the TGA stop codon are underlined.

SUBSTITUTE SHEET

20/26

Figure 7




Plot of AT/ATGC in sequence pVE36, from base 800 to base 4000
computed using an interval of 50 bases.

REGION A : N-terminal deletion + modification

REGION B : internal modification


Figure 7 (cont.)



Plot of AT/ATGC in sequence PPS029.
From base 800 to base 4000 computed using an interval of 50 bases.

22/26


FIGURE 8A



1 : AatII	8 : BclI	15 : EcoRV	22 : NarI	29 : RsrII
2 : AccI	9 : BspMII	16 : Eco31I	23 : NdeI	30 : SacI
3 : AflII	10 : BssHII	17 : EspI	24 : NheI	31 : SpeI
4 : AlwI	11 : BstEII	18 : HindII	25 : NsiI	32 : SphI
5 : ApaI	12 : BstXI	19 : HindIII	26 : PflMI	33 : Tth111I
6 : AvrII	13 : EagI	20 : KpnI	27 : PpuVI	34 : XcaI
7 : BanII	14 : EcoNI	21 : NaeI	28 : PstI	35 : XhoI


SUBSTITUTE SHEET

FIGURE 8A (CONT.)



1 : AatII	8 : BclI	15 : EcoRV	22 : NarI	29 : RsrII
2 : AccI	9 : BspMII	16 : Eco31I	23 : NdeI	30 : SacI
3 : AflII	10 : BssHII	17 : EspI	24 : NheI	31 : SpeI
4 : AlwNI	11 : BstEII	18 : HindII	25 : NsiI	32 : SphI
5 : ApaI	12 : BstXI	19 : HindIII	26 : PflMI	33 : Tth111I
6 : AvrII	13 : EagI	20 : KpnI	27 : PpuMI	34 : XcaI
7 : BamHI	14 : EcoNI	21 : NaeI	28 : PvuI	35 : XhoI

24/26


Figure 8b

SUBSTITUTE SHEET

25/26

Figure 8b (cont.)


TEX experiment construct PS029 in SAI protoplasts.



CIRCTITITE SHEET

Figure 8b(cont.)

TEX experiment construct PS029 in SFI protoplasts.
Bt2 Elisa 4/4/90 and CAT assay 5/4/90 .



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 91/00733

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/31 ; C12N15/82 ; C12N15/67 ; A01H5/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12N ; A01H	
Documentation Searched other than Minimum Documentation in the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	EP,A,359472 (LUBRIZOL GENETICS, INC.) 21 March 1990 see the whole document	1-3, 7, 9-17
Y	(cited in the application) ---	8
Y	NATURE. vol. 328, 02 July 1987, LONDON GB pages 33 - 37; VAECK, M. et al.: "Transgenic plants protected from insect attack" see the whole document (cited in the application) ---	8
A	EP,A,305275 (PLANT GENETIC SYSTEMS, N.V.) 01 March 1989 see the whole document --- -/--	9-17
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
04 JULY 1991	19 JUL 1991	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	ANDRES S.M.	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	PLANT PHYSIOLOGY vol. 85, 1987, pages 1103 - 1109; BARTON, K.A. et al.: "Bacillus thuringiensis delta-endotoxin expressed in transgenic Nicotiana tabacum provides resistance to Lepidopteran insects" see the whole document (cited in the application) ---	1-17
X,P	EP,A,385962 (MONSANTO COMPANY) 05 September 1990 see page 7, line 57 - page 12, line 30; figures 8, 9 see page 16, line 28 - page 22, line 39; claim 15 (cited in the application) ---	1-3, 7, 9-17

JA 40020

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
 The members are as contained in the European Patent Office EDP file on
 The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

04/07/91

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-359472	21-03-90	AU-A- 4118289	15-03-90
		JP-A- 2186989	23-07-90
EP-A-305275	01-03-89	AU-A- 2257288	09-03-89
		WO-A- 8901515	23-02-89
		JP-T- 2500566	01-03-90
EP-A-385962	05-09-90	AU-A- 5163090	26-09-90
		CA-A- 2024811	25-08-90
		EP-A- 0413019	20-02-91
		WO-A- 9010076	07-09-90

EPO FORM P0079

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82